

Citation for published version:

Pegg, E, Popat, B, Alinejad, M, Van Duren, B, Murray, DW & Pandit, HG 2014, 'In vivo knee kinematics of ACL-deficient patients after unicompartmental knee arthroplasty'.

Publication date:
2014

Document Version
Publisher's PDF, also known as Version of record

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

In vivo knee kinematics of ACL-deficient patients after unicompartmental knee arthroplasty



EC Pegg¹, B Popat¹, M Alinejad¹, BH van Duren¹, DW Murray¹, HG Pandit¹
¹ Nuffield Department of Orthopaedics, Rheumatology and Musculoskeletal Sciences (NDORMS), University of Oxford, UK


MAIN OBJECTIVE

- Compare the knee kinematics of ACL-deficient (ACLD) vs. ACL-intact (ACLI) patients after unicompartmental knee surgery using sagittal plane video fluoroscopy.

Measurements

The following measurements were manually made on each frame of every fluoroscopy video:

- Patellar Tendon Angle (PTA) = angle between axis 'C' and 'B'
- Knee Flexion Angle (KFA) = angle between 'A' and 'B'
- Bearing Movement (BM) = distance 'D'



METHODOLOGY


Patient selection

Case-control study where first ACLD patients were recruited prospectively, then ACLI patients were matched and then recruited for the control group. All patients had undergone Oxford unicompartmental knee replacement on the medial side of their knee between January 2000 and June 2011. This study was granted ethical approval in January 2013. A summary of the ACLD and ACLI cohort groups are shown below:


	ACLD	ACLI	p-value	Significance
# Knees	16	16		
# Patients	14	13		
Age	67.0 (50-87)	68.3 (49-86)	0.8046	NS
Follow-up Time	6.3 (1.3-12.8)	6.0 (2.6-11.0)	0.8209	NS
Gender	12 male, 2 female	12 male, 1 female	0.3173	NS
Oxford Knee Score	40.7 (20-48)	42.3 (32-48)	0.35	NS
ΔOxford Knee Score	15.9 (2-33)	12.9 (2-27)	0.57	NS
Tegner Activity Score	3.2 (2-5)	2.8 (0-5)	0.15	NS
VAS Pain Score	16.6 (0-70.3)	10.7 (0-85.9)	0.73	NS

Table 1. Description of the ACLD and ACLI patient cohorts. Data shown as: mean value (range). NS=not significant.

Step-up Exercise



Knee Bend Exercise



Knee fluoroscopy



All patients were asked to perform two exercises while a sagittal plane knee fluoroscopy was taken; a step-up exercise and a knee bend exercise (see images shown to the right).

Patients were allowed to stabilise themselves during the exercise using a handrail with their arm contralateral to the knee being examined. Patients were allowed one practise run for each exercise, after which the fluoroscopy was taken.


After each exercise a static image was taken of a calibration grid to ensure any pin-cushion or barrel distortion effects were removed. Distortion was removed using MATLAB software (version 7.1, MathWorks Inc. MA, USA) with a weighted least-squares method [1].

RESULTS

ACLD patients took ~3s longer to perform the exercises



Large anterior-posterior (AP) medial bearing movement observed in ACLD patients from 30-60 degrees of flexion



ACLD patients had a dip in PTA from 30-40 degrees of flexion

CONCLUSIONS

- Patients with ACL ligament deficiency after UKR have abnormal knee kinematics
- Differences were noticeable from 30-60 degrees of flexion and may relate to muscle imbalance [2]
- More variability was observed in AP bearing movement for ACLD patients
- The kinematics of ACLD-UKR knees were more normal than TKR, but less normal than ACLI-UKR and ACL-reconstructed-UKR knees [3].

ACKNOWLEDGEMENTS

- Biomet UK Healthcare Ltd.
- Jo Brown (NDORMS) and the hospital radiographers
- NIHR Biomedical Research Unit into Musculoskeletal Disease, NDORMS

[1] Goshtasby A. *Im. Vis. Comp.* 1988;6(4):255-61
[2] Berchuck M et al. *JBJS [Am]*. 1990;72(6):871-7.
[3] Pandit H et al. *JBJS [Br]*. 2005;87-B(7):940-5.